Author Index

Aderjan, R., see Bogusz, M. 43
Adinarayana, M.
——, Singh, U. S. and Dwivedi, T. S.
A biochromatographic technique for the quantitative estimation of triphenyltin fungicides 210
Aerts, M. M. L.
——, Beek, W. M. J. and Brinkman, U. A. Th.
Monitoring of veterinary drug residues by a combination of continuous flow techniques and column-switching high-performance liquid chromatography. I. Sulphonamides in egg, meat and milk using post-column derivatization with dimethylaminobenzaldehyde 97
Aguilar, M. I., see Hearn, M. T. W. 271
Ansari, G. A. S., see Bandi, Z. L. 513
Antia, F. D.
—— and Horváth, Cs.
High-performance liquid chromatography at elevated temperatures: examination of conditions for the rapid separation of large molecules 1
Bailly, J. R., see Leitao, J. 229
Bandi, Z. L.
—— and Ansari, G. A. S.
Separation of rat liver phospholipids from polar non-phospholipids by thin-layer chromatography 513
Beek, W. M. J., see Aerts, M. M. L. 97
Belenkii, B. G., see Tennikova, T. B. 357
Björkqvist, B.
—— and Hase, A.
Separation and determination of intact glucosinolates in rapeseed by high-performance liquid chromatography 501
Bogusz, M.
—— and Aderjan, R.
Improved standardization in reversed-phase high-performance liquid chromatography using 1-nitroalkanes as a retention index scale 43
Both-Miedema, R., see Jansen, E. H. J. M. 363
Brechbühler, B., see Schilling, B. 204
Brinkman, U. A. Th., see Aerts, M. M. L. 97
——, see Maris, F. A. 297
Burton, S. J.
——, McLoughlin, S. B., Stead, C. V. and Lowe, C. R.
Design and applications of biomimetic anthraquinone dyes. I. Synthesis and characterisation of terminal ring isomers of C.I. Reactive Blue 2 127
Cai, D.-G., see Zhang, T.-Y. 159
Chen, T.-S., see Philip, T. 113
Chhabra, B. R., see Dhillon, R. S. 256
Clark, M. A., see Strohsacker, M. W. 185
Coughlin, R. T., see Samu, J. 167
Čoupek, J., see Tennikova, T. B. 357
Crooke, S. T., see Strohsacker, M. W. 185
Dalene, M.
——, Mathiasson, L., Skarping, G., Sangö, C. and Sandström, J. F.
Trace analysis of airborne aromatic isocyanates and related aminoisocyanates and diamines using high-performance liquid chromatography with ultraviolet and electrochemical detection 469
——, see Skarping, G. 453
Damsté, J. S. Sinninghe, see Sinninghe Damsté, J. S. 435
De Jong, G. J., see Maris, F. A. 297
De Leeuw, J. W., see Sinninghe Damsté, J. S. 435
De Saint Blanquat, G., see Leitao, J. 229
Desta, B.
Complete separation of nine equine oestrogens by high-performance liquid chromatography 385
Dhillon, R. S.
——, Singh, J., Gautam, V. K. and Chhabra, B. R.
Detection of aminophenols and aromatic amines on thin-layer plates containing nitrite 256
Dohnal, J., see Jelínek, I. 496
Dorulla, G. K., see Riggin, R. M. 307
Dubs, R., see Schilling, B. 204
Dufek, P.
——, Vobecký, M., Holík, J. and Valášek, J.
Separation of some rare earths and transplutonium elements by high-performance liquid chromatography on Ostion cation exchanger 249
Duinker, J. S., see Petrick, G. 241
Dwivedi, T. S., see Adinarayana, M. 210
Elazhary, Y., see Mohanty, J. G. 149
Frei, R. W., see Maris, F. A. 297
Fridman, M., see Hearn, M. T. W. 271
Furukawa, M.
——, Yoshitake, N. and Yokoyama, T.
Identification of polymer glycols in polyurethane elastomers by pyrolysis–gas chromatography 219
Gautam, V. K., see Dhillon, R. S. 256
Gioeli, C., see Pettersson, C. 225
Goto, M., see Takeuchi, T. 482

Grob, K., see Schilling, B. 204
Gyllenhaal, O.
—— and Vessman, J.
Phosgene as a derivatizing reagent prior to gas and liquid chromatography (Review) 259
Hanaoka, Y., see Murayama, T. 417
Harvey, T. G.
——, Matheson, T. W. and Pratt, K. C.
Extraction and identification of nitrile compounds in rundle shale oil 193
Hase, A., see Björkqvist, B. 501
Hatano, H., see Murayama, T. 417
Hatano, T.
——, Yoshida, T. and Okuda, T.
Chromatography of tannins. III. Multiple peaks in high-performance liquid chromatography of some hydrolyzable tannins 285
Hearn, M. T. W.
——, Aguilar, M. I., Nguyen, T. and Fridman, M.
High-performance liquid chromatography of amino acids, peptides and proteins. LXXXIV. Application of derivative spectroscopy to the study of column residency effects in the reversed-phase and size-exclusion liquid chromatographic separation of proteins 271
Herche, L. R., see Parrish, C. C. 350
Holík, J., see Dufek, P. 249
Honda, A., see Ukai, S. 374
Horák, D., see Tennikova, T. B. 357
Horváth, Cs., see Antia, F. D. 1
——, see Velayudhan, A. 397
Hosoe, M.
——, Oi, T., Kawada, K. and Kakihana, H.
Separation of rubidium isotopes by ion-exchange chromatography 253
Hunter, K.
——, Sharp, E. A. and Newton, A.
Determination of diastereoisomers of bromadiolone, an anticoagulant rodenticide, in animal tissues by high-performance liquid chromatography 83
Ishii, D., see Takeuchi, T. 319, 482
Ito, Y., see Zhang, T.-Y. 159
Jansen, E. H. J. M.
—— and Both-Miedema, R.
Instability of phenylthiohydantoin amino acids 363
Jelínek, I.
——, Dohnal, J., Snopek, J. and Smolková-Keulemansová, E.
Use of cyclodextrins in isotachophoresis. III. Purity control of naftidrofuryl hydrogenoxalate and some of its synthesis intermediates 496
Jong, G. J. de, see Maris, F. A. 297
Kajiwara, M., see Kurumaya, K. 235
Kakihana, H., see Hosoe, P. 253
Kawada, K., see Hosoe, M. 253
Keler, T., see Samu, J. 167
Kennish, J. M., see Whitsett, J. F. 343
Kihara, K., see Murayama, T. 417
Kiho, T., see Ukai, S. 374
Kock-van Dalen, A. C., see Sinninghe Damsté, J. S. 435
Kolář, J., see Tennikova, T. B. 357
Kondo, N., see Takeuchi, T. 482
Kováts, E., see Samu, J. 167
Kubota, T., see Murayama, T. 417
Kurumaya, K.
——, Sakamoto, T., Okada, Y. and Kajiwara, M.
Application of droplet counter-current chromatography to the isolation of vitamin B_{12} 235
Kuwatsuka, S., see Spitzer, T. 489
Kwietniewski, L., see Wójcik, A. 55
Leeuw, J. W. de, see Sinninghe Damsté, J. S. 435
Lefeber, D. S., see Riggin, R. M. 307
Leitao, J.
——, De Saint Blanquat, G., Bailly, J. R. and Paillas, CH.
Quantitation of aflatoxins from various strains of aspergillus in foodstuffs 229
Lillig, B., see Velayudhan, A. 397
Liu, G.
——, Luo, X. and Zhang, S.
Determination of sulphur compounds at low parts per billion levels by gas chromatography with flame photometric detection 327
Lowe, C. R., see Burton, S. J. 127
Luo, X., see Liu, G. 327
McLoughlin, S. B., see Burton, S. J. 127
McNally, M. E. P.
—— and Wheeler, J. R.
Supercritical fluid extraction coupled with supercritical fluid chromatography for the separation of sulfonylurea herbicides and their metabolites from complex matrices 63
Maltzev, V. G., see Tennikova, T. B. 357
Maris, F. A.
——, Nijenhuis, M., Frei, R. W., De Jong, G. J. and Brinkman, U. A. Th.
On-line post-column extraction in column liquid chromatography with electron-capture detection. II. Applications 297
Martijn, A., see Schreuder, R. H. 73
Matheson, T. W., see Harvey, T. G. 193
Mathiasson, L., see Dalene, M. 469
——, see Skarping, G. 453
Miner, D. J., see Riggin, R. M. 307
Minnich, M. D., see Strohsacker, M. W. 185

Mohanty, J. G.
—— and Elazhary, Y.
High-performance liquid chromatographic separation and immunological characterization of soluble bovine viral diarrhea virus antigen 149
Mori, I., see Motohashi, N. 335
Moschidis, M. C.
Silicic acid column chromatography of phosphonolipids. IX. Silicic acid column chromatographic study of some phosphono analogues of bis(diacylglycero)phosphate and glycerophosphoric acid 508
Motohashi, N.
——, Nakamichi, Y., Mori, I., Nishikawa, H. and Umemoto, J.
Analysis by high-performance gel permeation chromatography of hyaluronic acid in animal skins and rabbit synovial fluid 335
Murayama, T.
——, Kubota, T., Hanaoka, Y., Rokushika, S., Kihara, K. and Hatano, H.
Peak enhancement of carboxylic acids in ion-exclusion chromatography with a cation-exchange hollow-fibre suppressor using neutral and alkaline enhancers 417
Nagai, K., see Ukai, S. 374
Nakamichi, Y., see Motohashi, N. 335
Nehmer, U.
Simultaneous determination of atracurium besylate and its major decomposition products and related impurities by reversed-phase high-performance liquid chromatography 425
Newton, A., see Hunter, K. 83
Nguyen, T., see Hearn, M. T. W. 271
Nguyen, V., see Samu, J. 167
Nijenhuis, M., see Maris, F. A. 297
Nishikawa, H., see Motohashi, N. 335
Noda, H.
——, Saitoh, K. and Suzuki, N.
Elution recovery of tris(acetylacetonato)aluminium(III), -gallium(III) and -indium(III) in size-exclusion chromatography 368
Nowotny, A., see Samu, J. 167
Nurmela, K. V. V.
—— and Satama, L. T.
Quantitative analysis of triglycerides by high-performance liquid chromatography using non-linear gradient elution and flame ionization detection 139
Oi, T., see Hosoe, M. 253
Okada, Y., see Kurumaya, K. 235
Okuda, T., see Hatano, T. 285
Paillas, CH., see Leitao, J. 229
Parrish, C. C.
——, Zhou, X. and Herche, L. R.
Flame ionization and flame thermionic detection of carbon and nitrogen in aquatic lipid and humic-type classes with an Iatroscan Mark IV 350
Persson, B., see Wannerberg, O. 199
Petrick, G.
——, Schulz, D. E. and Duinker, J. C.
Clean-up of environmental samples by high-performance liquid chromatography for analysis of organochlorine compounds by gas chromatography with electron-capture detection 241
Pettersson, C.
—— and Gioeli, C.
Improved resolution of enantiomers of naproxen by the simultaneous use of a chiral stationary phase and a chiral additive in the mobile phase 225
Philip, T.
—— and Chen, T.-S.
Separation and quantitative analysis of some carotenoid fatty acid esters on fruits by liquid chromatography 113
Pichler, P., see Schilling, B. 204
Poole, C. F., see Poole, S. K. 17
Poole, S. K.
—— and Poole, C. F.
Chemical interactions as a possible limitation on the useful solvent properties of liquid alkylammonium salts 17
Pratt, K. C., see Harvey, T. G. 193
Riggin, R. M.
——, Shaar, C. J., Dorulla, G. K., Lefeber, D. S. and Miner, D. J.
High-performance size-exclusion chromatographic determination of the potency of biosynthetic human growth hormone products 307
Rokushika, S., see Murayama, T. 417
Saint Blanquat, G. De, see Leitao, J. 229
Saitoh, K., see Noda, H. 368
Sakamoto, T., see Kurumaya, K. 235
Samu, J.
——, Kováts, E., Nguyen, V., Keler, T., Nowotny, A. and Coughlin, R. T.
Thin-layer chromatography of endotoxins, their derivatives and contaminants 167
Sandström, J. F., see Dalene, M. 469
Sangö, C., see Dalene, M. 469
Satama, L. T., see Nurmela, K. V. V. 139
Schenck, P. A., see Sinninghe Damsté, J. S. 435

Schilling, B.
——, Grob, K., Pichler, P., Dubs, R. and Brechbühler, B.
Fused-silica ends for glass capillaries: widening glass capillary butts by a heated tungsten point for preparing press-fit connections 204
Schreuder, R. H.
—— and Martijn, A.
Determination of dodecylbenzenesulphonates and ethoxylated alkylphenols in liquid pesticide formulations by high-performance liquid chromatography 73
Schulz, D. E., see Petrick, G. 241
Shaar, C. J., see Riggin, R. M. 307
Sharp, E. A., see Hunter, K. 83
Shorr, R. G. L., see Strohsacker, M. W. 185
Singh, J., see Dhillon, R. S. 256
Singh, U. S., see Adinarayana, M. 210
Sinninghe Damsté, J. S.
——, Kock-van Dalen, A. C., De Leeuw, J. W. and Schenck, P. A.
Identification of homologous series of alkylated thiophenes, thiolanes, thianes and benzothiophenes present in pyrolysates of sulphur-rich kerogens 435
Skarping, G., see Dalene, M. 469
——, Dalene, M. and Mathiasson, L.
Trace analysis of airborne 1,6-hexamethylene-diisocyanate and the related aminoisocyanate and diamine by glass capillary gas chromatography 453
Smolková-Keulemansová, E., see Jelínek, I. 496
Snopek, J., see Jelínek, I. 496
Spitzer, T.
—— and Kuwatsuka, S.
Simultaneous clean-up of nitroarenes and polycyclic aromatic ketones from soil and particulate matter on XAD-2 489
Stead, C. V., see Burton, S. J. 127
Strohsacker, M. W.
——, Minnich, M. D., Clark, M. A., Shorr, R. G. L. and Crooke, S. T.
Selective purification of cardiac myosin by a high-performance salicylate affinity column 185
Suzuki, N., see Noda, H. 368
Švec, F., see Tennikova, T. B. 357
Szabó-Ravasz, B.
High-performance liquid chromatographic methods for the determination of contaminants in technical phosmethylan 380
Takeuchi, T.
—— and Ishii, D.
Signal enhancement by on-column fluorometric detection in high-performance liquid chromatography using micropacked fused-silica columns 319
——, Watanabe, S., Kondo, N., Ishii, D. and Goto, M.
Improvement of the interface for coupling of fast atom bombardment mass spectrometry and micro high-performance liquid chromatography 482
Tennikova, T. B.
——, Horák, D., Švec, F., Kolář, J., Čoupek, J., Trushin, S. A., Maltzev, V. G. and Belenkii, B. G.
Hydrolysed macroporous glycidyl methacrylate–ethylene dimethacrylate copolymer sorbent for size-exclusion high-performance liquid chromatography of synthetic polymers and biopolymers 357
Trushin, S. A., see Tennikova, T. B. 357
Tsuchiya, T., see Ukai, S. 374
Ukai, S.
——, Honda, A., Nagai, K., Kiho, T. and Tsuchiya, T.
Gas–liquid chromatography of acid hydrolysates of hydroxyethylstarch and hydroxyethylcellulose 374
Umemoto, J., see Motohashi, N. 335
Valášek, J., see Dufek, P. 249
Velayudhan, A.
——, Lillig, B. and Horváth, Cs.
Analysis of multiple front formation in the wetting of thin-layer plates 397
Vessman, J., see Gyllenhaal, O. 259
Vobecký, M., see Dufek, P. 249
Voelkel, A.
Dispersive force parameters as a measure of the polarity of surfactants used as stationary phases in gas–liquid chromatography 29
Wannerberg, O.
—— and Persson, B.
Liquid chromatographic method for the determination of bambuterol hydrochloride and related compounds 199
Watanabe, S., see Takeuchi, T. 482
Wheeler, J. R., see McNally, M. E. P. 63
Whitsett, J. F.
—— and Kennish, J. M.
Calibration of thin-layer chromatography with flame ionization detection for the analysis of natural lipid samples 343
Wójcik, A.
—— and Kwietniewski, L.
Preparation of porous bead silica 55
Yokoyama, T., see Furukawa, M. 219
Yoshida, T., see Hatano, T. 285
Yoshitake, N., see Furukawa, M. 219
Zhang, S., see Liu, G. 327
Zhang, T.-Y.
——, Cai, D.-G. and Ito, Y.
Separations of flavonoids and alkaloids in medicinal herbs by high-speed counter-current chromatography 159
Zhou, X., see Parrish, C. C. 350